arXiv:1602.03569v1 [math.CO] 10 Feb 2016

THE INDEPENDENCE NUMBER OF NON-UNIFORM UNCROWDED HYPERGRAPHS AND AN ANTI-RAMSEY TYPE RESULT

SANG JUNE LEE AND HANNO LEFMANN

Abstract. We prove the following: Fix an integer $k \geq 2$, and let T be a real number with $T \geq 1.5$. Let $\mathcal{H} = (V, \mathcal{E}_2 \cup \mathcal{E}_3 \cup \cdots \cup \mathcal{E}_k)$ be a non-uniform hypergraph with the vertex set V and the set $\mathcal{E}_i$ of edges of size $i = 2, \ldots, k$. Suppose that $\mathcal{H}$ has no 2-cycles (regardless of sizes of edges), and neither contains 3-cycles nor 4-cycles consisting of 2-element edges. If the average degrees $t_i^{i-1} := i|\mathcal{E}_i|/|V|$ satisfy that $t_i^{i-1} \leq T^{i-1}(\ln T)^{\frac{k-i}{k-1}}$ for $i = 2, \ldots, k$, then there exists a constant $C_k > 0$, depending only on k, such that $\alpha(\mathcal{H}) \geq C_k \frac{|V|}{T}(\ln T)^{\frac{1}{k-1}}$, where $\alpha(\mathcal{H})$ denotes the independence number of $\mathcal{H}$. This extends results of Ajtai, Komlós, Pintz, Spencer and Szemerédi [J. Comb. Theory Ser. A 32, 1982, 321–335] and Duke, Rödl and the second author [Random Struct. Algorithms 6, 1995, 209–212] for uniform hypergraphs.

As an application, we consider an anti-Ramsey type problem on non-uniform hypergraphs. Let $\mathcal{H} = \mathcal{H}(n; 2, \ldots, \ell)$ be the hypergraph on the n-vertex set V in which, for $s = 2, \ldots, \ell$, each s-subset of V is a hyperedge of $\mathcal{H}$. Let Δ be an edge-coloring of $\mathcal{H}$ satisfying the following: (a) two hyperedges sharing a vertex have different colors; (b) two hyperedges with distinct size have different colors; (c) a color used for a hyperedge of size s appears at most u_s times. For such a coloring Δ, let $f_\Delta(n; u_2, \ldots, u_\ell)$ be the maximum size of a subset U of V such that each hyperedge of $\mathcal{H}[U]$ has a distinct color, and let $f(n; u_2, \ldots, u_\ell) := \min_\Delta f_\Delta(n; u_2, \ldots, u_\ell)$. We determine $f(n; u_2, \ldots, u_\ell)$ up to a multiplicative logarithm factor, which is a non-uniform version of a result for edge-colorings of graphs by Babai [Graphs Comb. 1, 1985, 23–28], and for uniform hypergraphs by Alon, Rödl and the second author [Coll. Math. Soc. János Bolyai, 60. Sets, Graphs and Numbers, 1991, 9–22] and by Rödl, Wysocka and the second author [J. Comb. Theory Ser. A 74, 1996, 209–248].

1. Introduction

Let $\mathcal{H} = (V, \mathcal{E})$ be a hypergraph with its vertex set V and its edge set $\mathcal{E}$. Let $\mathcal{E}_i \subset \mathcal{E}$ be the set of all i-element edges in $\mathcal{H}$. For a vertex $v \in V$, let $d_i(v)$ denote the number of i-element edges $E \in \mathcal{E}_i$ containing v. A hypergraph $\mathcal{H} = (V, \mathcal{E})$ is called *k-uniform* if $\mathcal{E} = \mathcal{E}_k$. A subset $V' \subseteq V$ is called *independent* if for no edge $E \in \mathcal{E}$ it is $E \subseteq V'$. The *independence number* $\alpha(\mathcal{H})$ of $\mathcal{H}$ is the maximum size of an independent set of $\mathcal{H}$. For a subset $V^* \subset V$ of the vertex set, let $\mathcal{H}[V^*]$ be the subhypergraph of $\mathcal{H}$ induced on V^*.

The independence number has been well-studied for uniform hypergraphs, however, for non-uniform hypergraphs it was not that studied correspondingly. The goal of this paper is to extend known results on the independence number of a uniform hypergraph to a non-uniform hypergraph.

Turán's theorems by Turán [8] and Spencer [9] imply the following theorem about the independence number of a k-uniform hypergraph $\mathcal{H}$.

Theorem 1 (Turán [8] and Spencer [9])**.** *Let $k \geq 2$ be an integer. Let $\mathcal{H} = (V, \mathcal{E}_k)$ be a k-uniform hypergraph with N vertices and average degree $t^{k-1} := k|\mathcal{E}_k|/N$, where $t \geq 1$. Then,*

$$\alpha(\mathcal{H}) \geq \frac{k-1}{k}\frac{N}{t}\,. \tag{1}$$

Date: Fri 12th Feb, 2016, 1:24am.

Later, a better lower bound on the independence number of a k-uniform hypergraph was obtained if the hypergraph does not contain cycles of small lengths. We introduce the definition of cycles of a given length.

Definition 2. *Let $\mathcal{H} = (V, \mathcal{E})$ be a hypergraph. A j-*cycle *in $\mathcal{H}$ is a family of pairwise distinct edges $E_1, \ldots, E_j \in \mathcal{E}$ such that the following hold:*

- *$E_i \cap E_{i+1} \neq \emptyset$ for $i = 1, \ldots, j-1$ and $E_j \cap E_1 \neq \emptyset$.*
- *There are pairwise distinct vertices $v_1, \ldots, v_j$ such that $v_i \in E_i \cap E_{i+1}$ for $i = 1, \ldots, j-1$ and $v_j \in E_j \cap E_1$.*

Definition 3. *A hypergraph $\mathcal{H}$ is called* uncrowded *if it does not contain any 2-, 3- or 4-cycles. A hypergraph $\mathcal{H}$ is called* linear *if it does not contain any 2-cycles.*

Ajtai, Komlós, Pintz, Spencer and Szemerédi [1] obtained a lower bound on the independence number of a k-uniform uncrowded hypergraph as follows. Later, Bertram-Kretzberg and Lefmann [4] and Fundia [6] provided a deterministic polynomial time algorithm.

Theorem 4 (Ajtai, Komlós, Pintz, Spencer and Szemerédi [1])**.** *Let $k \geq 2$ be a fixed integer. Let t and N satisfy $t > t_0(k)$ and $N > N_0(k, t)$. Let $\mathcal{H} = (V, \mathcal{E}_k)$ be an uncrowded k-uniform hypergraph on N vertices with average degree $t^{k-1} := k|\mathcal{E}_k|/N$. Then, there exists a constant $C_k > 0$ such that*

$$\alpha(\mathcal{H}) \geq C_k \frac{N}{t} (\ln t)^{\frac{1}{k-1}} \ . \tag{2}$$

We remark that in [1] the constant C_k is bounded from below by $C_k \geq \frac{0.98}{e} 10^{-\frac{5}{k-1}}$.

Duke, Lefmann, and Rödl in [5] weakened the assumption in Theorem 4 for $k \geq 3$ as follows:

Theorem 5 (Duke, Lefmann, and Rödl [5])**.** *Let $k \geq 3$ be a fixed integer. Let t and N satisfy $t > t_0(k)$ and $N > N_0(k, t)$. Let $\mathcal{H} = (V, \mathcal{E}_k)$ be a linear k-uniform hypergraph on N vertices with average degree $t^{k-1} := k|\mathcal{E}_k|/N$. Then, there exists a constant $C'_k > 0$ such that*

$$\alpha(\mathcal{H}) \geq C'_k \frac{N}{t} (\ln t)^{\frac{1}{k-1}} \ . \tag{3}$$

In this paper we extend Theorems 4 and 5 to non-uniform hypergraphs as follows.

Theorem 6. *Let $k \geq 2$ be a fixed integer. Let T be a real number with $T \geq 1.5$ and N be a positive integer. Let $\mathcal{H} = (V, \mathcal{E}_2 \cup \cdots \cup \mathcal{E}_k)$ be a linear hypergraph on N vertices such that there are no 3-cycles and 4-cycles consisting of 2-element edges. Let the average degrees $t_i^{i-1} := i|\mathcal{E}_i|/N$ satisfy that for $i = 2, \ldots, k$*

$$t_i^{i-1} \leq T^{i-1} (\ln T)^{\frac{k-i}{k-1}}. \tag{4}$$

Then, there exists a constant $C_k > 0$ such that

$$\alpha(\mathcal{H}) \geq C_k \frac{N}{T} (\ln T)^{\frac{1}{k-1}} \ . \tag{5}$$

Remark 7. *For the range of $0 < T < 1.5$ in Theorem 6, a simple greedy algorithm gives that there exists a constant $C'_k > 0$, depending only on k, such that $\alpha(\mathcal{H}) \geq C'_k N$.*

As an application of the main theorem (Theorem 6), we consider a anti-Ramsey type problem of a non-uniform hypergraph. Let $\mathcal{H} = \mathcal{H}(n; 2, \ldots, \ell)$ be the hypergraph on the vertex set V, $|V| = n$, in which, for $s = 2, \ldots, \ell$, each s-subset of V is a hyperedge of $\mathcal{H}$. Suppose that Δ is an edge-coloring of $\mathcal{H}$ satisfying the following conditions:

(a) Two hyperedges sharing a vertex have different colors. In other words, each color class is a *matching*.

(b) Two hyperedges with distinct size have different colors.

(c) The coloring is $(u_2, \ldots, u_\ell)$-*bounded*, that is, a color used for a hyperedge of size s appears at most u_s times.

For such a coloring Δ, let $f_\Delta(n; u_2, \ldots, u_\ell)$ be the maximum size of a vertex set U in V such that the subhypergraph $\mathcal{H}[U]$ of $\mathcal{H}$ induced on U is *totally multicolored*, which means that each hyperedge of $\mathcal{H}[U]$ has a distinct color. Let

$$f(n; u_2, \ldots, u_\ell) := \min_\Delta f_\Delta(n; u_2, \ldots, u_\ell),$$

where we minimize it over all edge-colorings Δ of $\mathcal{H}$ satisfying (a)–(c).

We will show the following which is a non-uniform version of a result for edge-colorings of graphs in [7].

Theorem 8.

(a) *Suppose that*

$$\max_{2\le i\le \ell}\left[\left(n^{i-1}u_i\right)^{\frac{1}{2i-1}}\right] = \left(n^{s-1}u_s\right)^{\frac{1}{2s-1}}. \tag{6}$$

Then, there exist positive constants c_1 and c_2, depending only on ℓ, such that for every sufficiently large n,

$$c_1\left(\frac{n^s}{u_s}\right)^{\frac{1}{2s-1}} \le f(n; u_2, ..., u_\ell) \le c_2\left(\frac{n^s}{u_s}\ln n\right)^{\frac{1}{2s-1}}. \tag{7}$$

(b) *Suppose that*

$$\max_{2\le i\le \ell}\left[\left(\frac{n^{i-1}u_i}{\ln n}\right)^{\frac{1}{2i-1}}\right] = \left(\frac{n^{s-1}u_s}{\ln n}\right)^{\frac{1}{2s-1}} \tag{8}$$

and

$$u_s \ge n^{1/2+\varepsilon} \qquad \text{for some absolute constant } \varepsilon > 0. \tag{9}$$

Then, there exist positive constants $c_3 = c_3(\ell, \varepsilon)$ and $c_4 = c_4(\ell)$ such that for every sufficiently large n,

$$c_3\left(\frac{n^s}{u_s}\ln n\right)^{\frac{1}{2s-1}} \le f(n; u_2, ..., u_\ell) \le c_4\left(\frac{n^s}{u_s}\ln n\right)^{\frac{1}{2s-1}}. \tag{10}$$

Remark 9. *Statements (a) and (b) in Theorem 8 have different assumptions. We remark that under asumption* (8)*, our argument for the lower bound in* (7) *gives*

$$c_3'\left(\frac{n^s}{u_s}\right)^{\frac{1}{2s-1}}(\ln n)^{\frac{1}{2s-1}-\frac{1}{2\ell-1}} \le f(n; u_2, ..., u_\ell),$$

which is less than the lower bound in (10)*, while under assumption* (6)*, our argument for the lower bound in* (10) *gives*

$$c_3''\left(\frac{n^s}{u_s}\right)^{\frac{1}{2s-1}}\max\left[1, \left(\ln\frac{u_s^2}{n}\right)^{\frac{1}{2\ell-1}}\right] \le f(n; u_2, ..., u_\ell),$$

which is bigger than the lower bound in (7) *if $u_s \gg \sqrt{n}$.*

If we assume $u_2 = \ldots = u_\ell = n$, Theorem 8 (b) immediately implies the following corollary:

Corollary 10. *There exist positive constants c_3 and c_4, depending only on ℓ, such that for every sufficiently large n,*

$$c_3\left(n\ln n\right)^{\frac{1}{3}} \le f(n; \underbrace{n, ..., n}_{\ell-1}) \le c_4\left(n\ln n\right)^{\frac{1}{3}}. \tag{11}$$

For the special case of $\ell = 2$, i.e., edge-coloings of graphs, the upper bound in (11) was shown in 1985 by Babai [3], and the lower bound was proved in 1991 by Alon, Rödl and the second author [2].

The organization of this paper is as follows. Section 2 contains an extension of Theorem 1 to non-uniform hypergraphs. In Section 3, we show an extension of Theorem 4 to non-uniform hypergraphs. Section 4 contains our proof of an extension of Theorem 5 to non-uniform hypergraphs. In Section 5, we show Theorem 6 in full. Section 6 contains the proof of Theorem 8.

2. Arbitrary non-uniform hypergraphs

The next theorem by Spencer [9] provides a lower bound on the independence number $\alpha(\mathcal{H})$ of a non-uniform hypergraph $\mathcal{H}$.

Theorem 11 (Spencer [9])**.** *Let $\mathcal{H} = (V, \mathcal{E}_2 \cup \cdots \cup \mathcal{E}_k)$ be a hypergraph with N vertices and average degree $t_i^{i-1} := i|\mathcal{E}_i|/N$ for the i-element edges, where $i = 2, \ldots, k$. Let $T := \max\{t_i \mid 2 \le i \le k\} \ge 1/2$. Then,*

$$\alpha(\mathcal{H}) \ge \frac{1}{4}\frac{N}{T}.$$

Proof. For convenience we provide the sketch of a proof. Choose each vertex uniformly at random and independently with probability $p := 1/(2T) \le 1$. Let V^* be the random set of chosen vertices and let $\mathcal{E}_i^*$, $i = 2, \ldots, k$, be the sets of i-element edges in the induced random subhypergraph $\mathcal{H}[V^*]$. Then we have in expectation

$$\begin{aligned}\mathbb{E}\left[|V^*| - \sum_{i=2}^{k} |\mathcal{E}_i^*|\right] &= \mathbb{E}[|V^*|] - \sum_{i=2}^{k} \mathbb{E}[|\mathcal{E}_i^*|] = pN - \sum_{i=2}^{k} \frac{p^i N t_i^{i-1}}{i} \\ &\ge pN - \sum_{i=2}^{k} \frac{p^i N T^{i-1}}{i} = \frac{N}{2T} - \sum_{i=2}^{k} \frac{1}{i2^i}\frac{N}{T} \ge \frac{N}{4T}.\end{aligned}$$

Thus there exists a subset $V^* \subseteq V$ such that

$$|V^*| - \sum_{i=2}^{k} |\mathcal{E}_i^*| \ge \frac{N}{4T}.$$

By deleting one vertex from each edge $E \in \mathcal{E}_i^*$, $i = 2, \ldots, k$, we destroy all edges in $\mathcal{H}[V^*]$, and hence we obtain an independent set V^{**} of $\mathcal{H}$ with $|V^{**}| \ge N/(4T)$. □

3. Uncrowded non-uniform hypergraphs

In this section we show the following lemma which gives a lower bound on the independence number of an uncrowded non-uniform hypergraph.

Lemma 12. *Let $k \ge 2$ be a fixed integer. Let T and N satisfy $T > T_0(k)$ and $N > N_0(k, T)$. Let $\mathcal{H} = (V, \mathcal{E}_2 \cup \cdots \cup \mathcal{E}_k)$ be an uncrowded hypergraph on N vertices with average degrees $t_i^{i-1} := i|\mathcal{E}_i|/N$ such that for $i = 2, \ldots, k$*

$$t_i^{i-1} \le c_i T^{i-1} (\ln T)^{\frac{k-i}{k-1}},$$

where c_i are constants satisfying $0 < c_i < \frac{1}{16k^2}\binom{k-1}{i-1} 10^{-3\frac{(k-i)}{k-1}}$. Then, there exists a constant $C_k > 0$ such that

$$\alpha(\mathcal{H}) \ge C_k \frac{N}{T} (\ln T)^{\frac{1}{k-1}}. \tag{12}$$

In order to prove Lemma 12, we will use the following lemma which was used to show Theorem 4 in Ajtai, Komlós, Pintz, Spencer and Szemerédi [1]. Since the lemma in [1] has been written densely (with several typos), we provide a slightly modified statement, based on Lemma 6 in Fundia [6].

Lemma 13 (Ajtai, Komlós, Pintz, Spencer and Szemerédi [1])**.** *Let $k \geq 2$ be a fixed integer. Let T, N, and s be such that*

$$T \geq T_0(k), \;\; N \geq N_0(k,T), \;\; 0 \leq s \leq 0.01 \ln T. \tag{13}$$

Let n and t be integers satisfying

$$\frac{1}{2}\frac{N}{e^s} \leq n \leq \frac{N}{e^s} \quad \text{and} \quad \frac{T}{e^s} \leq t \leq 2\frac{T}{e^s}. \tag{14}$$

Let $\mathcal{H} = (V, \mathcal{E}_2 \cup \cdots \cup \mathcal{E}_k)$ be an uncrowded hypergraph with n vertices satisfying the following: for each vertex $v \in V$ and $2 \leq i \leq k$, the numbers $d_i(v)$ of i-element edges $E \in \mathcal{E}_i$ containing vertex v fulfill

$$d_i(v) \leq \binom{k-1}{i-1} s^{\frac{k-i}{k-1}} t^{i-1}.$$

Then, there exists an independent set $I \subseteq V$ and a vertex set $V^* \subset V$ with $I \cap V^* = \emptyset$ satisfying the following:

(i) $\alpha(\mathcal{H}) \geq |I| + \alpha(\mathcal{H}^*)$ *where* $\mathcal{H}^* = \mathcal{H}[V^*]$

(ii) $|I| \geq \dfrac{0.99}{e} w_s \dfrac{n}{t}$ *where* $w_s := (s+1)^{\frac{1}{k-1}} - s^{\frac{1}{k-1}}$

(iii) $\dfrac{n}{e}(1-\varepsilon) \leq |V^*| \leq \dfrac{n}{e}$ *where* $\varepsilon = 1/(10^6 \ln T)$

(iv) *for every vertex $v \in V^*$ and $i = 2, \ldots, k$ it is*

$$d_i^*(v) \leq \binom{k-1}{i-1} (s+1)^{\frac{k-i}{k-1}} \left(\frac{t}{e}(1+\varepsilon)\right)^{i-1},$$

where $d_i^*(v)$ is the number of i-element edges in $\mathcal{H}^*$ containing v.

In order to apply Lemma 13, we are going to obtain an induced subhypergraph of $\mathcal{H}$ in Lemma 12 that satisfies the assumptions of Lemma 13.

Lemma 14. *Let $k \geq 2$ be a fixed integer. Let T, N and s be positive integers satisfying*

$$T \geq T_0(k), \;\; N \geq N_0(k,T), \;\; s = 0.001 \ln T.$$

Let $\mathcal{H} = (V, \mathcal{E}_2 \cup \cdots \cup \mathcal{E}_k)$ be a hypergraph with N vertices and average degrees $t_i^{i-1} := i|\mathcal{E}_i|/N$ satisfying that

$$t_i^{i-1} \leq c_i T^{i-1} (\ln T)^{\frac{k-i}{k-1}},$$

where the constants $c_i > 0$ are such that $c_i < \frac{1}{16k^2}\binom{k-1}{i-1} 10^{-3\frac{k-i}{k-1}}$.

Then, $\mathcal{H}$ contains a subhypergraph $\mathcal{H}^* = (V^*, \mathcal{E}_2^* \cup \cdots \cup \mathcal{E}_k^*)$ induced on V^* with $|V^*| = n$ such that the following hold.

(a) $\dfrac{3}{4}\dfrac{N}{e^s} \leq n \leq \dfrac{N}{e^s}$, *and*

(b) *for each vertex $v \in V^*$,*

$$d_i^*(v) \leq \binom{k-1}{i-1} s^{\frac{k-i}{k-1}} \left(\frac{T}{e^s}\right)^{i-1}, \tag{15}$$

where $d_i^*(v)$ is the number of i-element edges in $\mathcal{H}^*$ containing v.

Proof. Let V' be a random set obtained by choosing each vertex in $\mathcal{H}$ independently with probability $p := 1/e^s$. For its expected size we have $\mathbb{E}[|V'|] = Np = N/e^s$, thus by Chernoff's inequality it is

$$\Pr\big(|V'| \leq \mathbb{E}[|V'|] - u\big) \leq e^{-u^2/N}$$

for every real $u \geq 0$. Then, with $s = 10^{-3} \ln T$ and a sufficiently large $N \geq N_0(k,T)$ we have

$$\Pr\left(|V'| \leq \mathbb{E}[|V'|] - \frac{1}{8}\frac{N}{e^s}\right) \leq e^{-\frac{N^2/(64e^{2s})}{N}} = e^{-N/(64e^{2s})} < \frac{1}{k}. \tag{16}$$

Let $\mathcal{H}' = (V', \mathcal{E}'_2 \cup \cdots \cup \mathcal{E}'_k)$ be the subhypergraph of $\mathcal{H}$ induced on V'. For $i = 2, \ldots, k$, we have

$$\mathbb{E}[|\mathcal{E}'_i|] = p^i |\mathcal{E}_i| = p^i t_i^{i-1} N/i \leq e^{-si} c_i T^{i-1} (\ln T)^{\frac{k-i}{k-1}} N/i.$$

By Markov's inequality we infer

$$\Pr\left(|\mathcal{E}'_i| > k\mathbb{E}[|\mathcal{E}'_i|]\right) \leq 1/k. \tag{17}$$

By (16) and (17), there exists an induced subhypergraph $\mathcal{H}' = (V', \mathcal{E}'_2 \cup \cdots \cup \mathcal{E}'_k)$ of $\mathcal{H}$ such that

$$|V'| \geq \frac{7}{8}\frac{N}{e^s}, \tag{18}$$

$$|\mathcal{E}'_i| \leq kc_i e^{-si} T^{i-1} (\ln T)^{\frac{k-i}{k-1}} N/i \quad \text{for } i = 2, \ldots, k. \tag{19}$$

From now on, we are going to obtain an induced subhypergraph $\mathcal{H}^*$ of $\mathcal{H}'$ such that the maximum degree is bounded. From (19), we infer

$$\mathbb{E}[d'_i(v)] \leq kc_i e^{-si} T^{i-1} (\ln T)^{\frac{k-i}{k-1}} N/|V'|.$$

Let Y_i be the number of vertices $v \in V'$ of high degree, i.e., such that

$$d'_i(v) > 7k \cdot kc_i e^{-si} T^{i-1} (\ln T)^{\frac{k-i}{k-1}} N/|V'|.$$

Since

$$Y_i \cdot 7k^2 c_i e^{-si} T^{i-1} (\ln T)^{\frac{k-i}{k-1}} N/|V'| \leq i|\mathcal{E}'_i| \leq kc_i e^{-si} T^{i-1} (\ln T)^{\frac{k-i}{k-1}} N,$$

we infer $Y_i \leq |V'|/(7k)$. Thus, the total number of vertices of high degree is bounded as

$$\sum_{i=2}^{k} Y_i \leq \sum_{i=2}^{k} \frac{|V'|}{7k} \leq \frac{|V'|}{7}. \tag{20}$$

By deleting these vertices of high degree from V', we obtain an induced subhypergraph $\mathcal{H}^* = (V^*, \mathcal{E}^*_2 \cup \cdots \cup \mathcal{E}^*_k)$ of $\mathcal{H}^*$ with $|V^*| \geq (6/7)|V'|$ such that with (18) we infer

$$\begin{aligned} |V^*| &\overset{(20)}{\geq} \frac{6}{7}|V'| \overset{(18)}{\geq} \frac{3}{4}\frac{N}{e^s} \\ d^*_i(v) &\leq 7k^2 c_i e^{-si} T^{i-1} (\ln T)^{\frac{k-i}{k-1}} N/|V'| \\ &\overset{(18)}{\leq} 8k^2 c_i \left(\frac{T}{e^s}\right)^{i-1} (\ln T)^{\frac{k-i}{k-1}}, \end{aligned} \tag{21}$$

where $d^*_i(v)$ is the number of i-element edges in $\mathcal{H}^*$ containing v. Recalling $s := 10^{-3} \ln T$ and the upper bound on c_i, we have

$$d^*_i(v) \leq \binom{k-1}{i-1} s^{\frac{k-i}{k-1}} \left(\frac{T}{e^s}\right)^{i-1},$$

which proves condition (b) in (15). We can obtain the condition $|V^*| \leq N/e^s$ by possibly deleting some more vertices from V^*. This together with (21) implies condition (a). □

Now we are ready to prove Lemma 12.

Proof of Lemma 12. We apply Lemma 14 with $s = 10^{-3} \ln T$ to $\mathcal{H}$ on N vertices. Then we obtain a subhypergraph $\mathcal{H}^* = (V^*, \mathcal{E}^*_2 \cup \cdots \cup \mathcal{E}^*_k)$ induced on V^* with $|V^*| = n$ such that the following hold:

(a) $\frac{3}{4}\frac{N}{e^s} \le n \le \frac{N}{e^s}$, and
(b) for each vertex $v \in V^*$, it is

$$d_i^*(v) \le \binom{k-1}{i-1} s^{\frac{k-i}{k-1}} \left(\frac{T}{e^s}\right)^{i-1},$$

where $d_i^*(v)$ is the number of i-element edges in $\mathcal{H}^*$ containing v.

Set $n_s = n$ and $t_s = T/e^s$ and $\mathcal{H}_s = \mathcal{H}^* = (V_s, \mathcal{E}_{2;s} \cup \cdots \cup \mathcal{E}_{k;s})$. We apply Lemma 13 with $\varepsilon = 1/(10^6 \ln T)$ to $\mathcal{H}_s$, iteratively. Let n_r and t_r be such that

$$\frac{3}{4}\frac{N}{e^r}(1-\varepsilon)^{r-s} \le n_r \le \frac{N}{e^r} \tag{22}$$

and

$$t_r = \frac{T}{e^r}(1+\varepsilon)^{r-s}. \tag{23}$$

For each $r = s, \ldots, 10^{-2} \ln T$, we obtain an independent set $I_r \subseteq V_r$ and a vertex set $V_{r+1} \subset V_r$ with $I_r \cap V_{r+1} = \emptyset$ satisfying the following:

(i) $\alpha(\mathcal{H}_r) \ge |I_r| + \alpha(\mathcal{H}_{r+1})$ where $\mathcal{H}_{r+1} = \mathcal{H}_r[V_{r+1}]$
(ii) $|I_r| \ge \frac{0.99}{e} w_r \frac{n_r}{t_r}$ where $w_r := (r+1)^{\frac{1}{k-1}} - r^{\frac{1}{k-1}}$
(iii) $\frac{n_r}{e}(1-\varepsilon) \le |V_{r+1}| \le \frac{n_r}{e}$ recalling $\varepsilon = 1/(10^6 \ln T)$
(iv) for every vertex $v \in V_{r+1}$ and $2 \le i \le k$,

$$d_{i;r+1}(v) \le \binom{k-1}{i-1} (s+1)^{\frac{k-i}{k-1}} (t_{r+1})^{i-1},$$

where $d_{i;r+1}(v)$ is the number of i-element edges in $\mathcal{H}_{r+1}$ containing v.

We first check how many times we can iteratively apply Lemma 13 to $\mathcal{H}_r$. Observe that we can apply Lemma 13 as far as inequalities (13) and (14) are satisfied.

- The inequality (13) is $s + r \le 0.01(\ln T)$.
- From (22), the inequalities about n in (14) are satisfied if $\frac{1}{2} \le \frac{3}{4}(1-\varepsilon)^{r-s}$. Using $1-p \ge e^{-2p}$ for $0 \le p \le 0.5$, one can check that it suffices to have $r \le 10^5 \ln T$.
- From (23), the inequalities about t in (14) are satisfied if $(1+\varepsilon)^{r-s} \le 2$. One can check that it suffices to have $r \le 10^5 \ln T$.

Therefore, we can apply Lemma 13 for $r + s \le 0.01 \ln T$.

Now we estimate the size of an independent set in $\mathcal{H}$ obtained by the above procedure. Notice that by using $(1+\varepsilon)^n \ge 1 + \varepsilon n$ and $1 + \varepsilon \le e^{\varepsilon}$ and $r \le 10^{-2} \ln T$ and $\varepsilon = 10^{-6}/\ln T$ we have

$$\begin{aligned} \frac{n_r}{t_r} &\ge \frac{(3/4)N(1-\varepsilon)^{r-s}/e^r}{T(1+\varepsilon)^{r-s}/e^r} \ge \frac{3}{4}\frac{N}{T}\frac{(1-\varepsilon)^r}{(1+\varepsilon)^r} \\ &\ge \frac{3}{4}\frac{N}{T}\frac{1-\varepsilon r}{e^{\varepsilon r}} \ge \frac{3}{4}\frac{N}{T}\frac{1-10^{-8}}{e^{10^{-8}}} \ge 0.74\frac{N}{T}. \end{aligned}$$

Hence, recalling that $w_r = (r+1)^{\frac{1}{k-1}} - r^{\frac{1}{k-1}}$, we obtain an independent set $I = I_s \cup \cdots \cup I_{0.01 \ln T}$ in $\mathcal{H}$ with

$$\begin{aligned}\alpha(\mathcal{H}) &\geq |I| = \sum_{r=s}^{0.01\ln T} |I_r| \geq \frac{0.99}{e} 0.74 \frac{N}{T} \sum_{r=s}^{0.01\ln T} w_r \\ &\geq \frac{0.73}{e}\frac{N}{T} \sum_{r=s}^{0.01\ln T} \left((r+1)^{\frac{1}{k-1}} - r^{\frac{1}{k-1}} \right) \\ &\geq \frac{0.73}{e}\frac{N}{T} (\ln T)^{\frac{1}{k-1}} \left(0.01^{\frac{1}{k-1}} - 0.001^{\frac{1}{k-1}} \right),\end{aligned}$$

which gives the lower bound (12) in Lemma 12. □

4. A weak version of Theorem 6

Now we show a (seemingly) weaker version of Theorem 6 in which T and N are large and the assumptions of the upper bounds on t_i^{i-1} are a bit different.

Lemma 15. *Let $k \geq 2$ be a fixed integer. Let T and N satisfy $T > T_0(k)$ and $N > N_0(k,T)$. Let $\mathcal{H} = (V, \mathcal{E}_2 \cup \cdots \cup \mathcal{E}_k)$ be a linear hypergraph on N vertices such that there are no 3-cycles and 4-cycles consisting of 2-element edges. Let the average degrees $t_i^{i-1} := i|\mathcal{E}_i|/N$ satisfy for $i = 2, \ldots, k$*

$$t_i^{i-1} \leq c_i T^{i-1} (\ln T)^{\frac{k-i}{k-1}}, \tag{24}$$

where c_i are constants satisfying $0 < c_i < \frac{1}{2^9 k^6} \binom{k-1}{i-1} 10^{-3\frac{k-i}{k-1}}$. Then, there exists a constant $C_k > 0$ such that

$$\alpha(\mathcal{H}) \geq C_k \frac{N}{T} (\ln T)^{\frac{1}{k-1}} . \tag{25}$$

Proof. Let $d_i(v)$ denote the number of i-element edges in $\mathcal{H}$ containing vertex $v \in V$. We delete all vertices $v \in V$ with $d_i(v) > k c_i T^{i-1} (\ln T)^{\frac{k-i}{k-1}}$ for some $i = 2, \ldots, k$. Let V^* be the set of remaining vertices, hence we have $|V^*| \geq N/k$. Then, the subhypergraph $\mathcal{H}^* = (V^*, \mathcal{E}_2^* \cup \cdots \cup \mathcal{E}_k^*)$ of $\mathcal{H}$ induced on V^* satisfies that for each vertex $v \in V^*$

$$d_i^*(v) \leq k c_i T^{i-1} (\ln T)^{\frac{k-i}{k-1}}, \tag{26}$$

where $d_i^*(v)$ is the number of i-element edges in $\mathcal{H}^*$ containing vertex v.

Now we estimate the numbers of 3-cycles and of 4-cycles not containing a 3-cycle in $\mathcal{H}^*$. First we consider the number of 3-cycles in $\mathcal{H}^*$. Let $C^*(g,h,i)$ denote the number of 3-cycles $\{E_1, E_2, E_3\}$ in $\mathcal{H}^*$ such that $|E_1| = g$, $|E_2| = h$, and $|E_3| = i$. To estimate $C^*(g,h,i)$, we fix a vertex $v \in V^*$ and regard it as a vertex in $E_1 \cap E_2$. There are at most $k c_g T^{g-1} (\ln T)^{\frac{k-g}{k-1}}$ edges in $\mathcal{E}_g^*$ containing vertex v. Let E_1 be one of these edges. Similarly, there are at most $k c_h T^{h-1} (\ln T)^{\frac{k-h}{k-1}}$ edges in $\mathcal{E}_h^*$ containing v. Let E_2 be one of these edges. Moreover, we fix one vertex $w \in E_1$ with $w \neq v$ and another vertex $x \in E_2$ with $x \neq v$ in at most $(k-1)^2$ ways, and consider $v, x \in E_3$. Since $\mathcal{H}^*$ is linear, there is at most one i-element edge $E_3 \in \mathcal{E}_i^*$ containing both vertices w and x. Hence, for each $2 \leq g \leq h \leq i \leq k$,

$$\begin{aligned}C^*(g,h,i) &\leq N \cdot k c_g T^{g-1} (\ln T)^{\frac{k-g}{k-1}} \cdot k c_h T^{h-1} (\ln T)^{\frac{k-h}{k-1}} \cdot (k-1)^2 \\ &< N k^4 c_g c_h T^{g+h-2} (\ln T)^2 .\end{aligned} \tag{27}$$

Next, we consider the number of 4-cycles in $\mathcal{H}^*$. Let $C^*(g,h,i,j)$ be the number of 4-cycles $\{E_1, E_2, E_3, E_4\}$ in $\mathcal{H}^*$ such that $|E_1| = g$, $|E_2| = h$, $|E_3| = i$, and $|E_4| = j$ and any three of

$\{E_1, E_2, E_3, E_4\}$ do not form a 3-cycle. With an argument similar to the argument to obtain (27), one can show that for each $2 \le g \le h \le i \le j \le k$,

$$C^*(g,h,i,j) < 3Nk^6 c_g c_h c_i T^{g+h+i-3} (\ln T)^3. \tag{28}$$

Note that the factor 3 in (28) arises due to the possible arrangements of the edges of possibly different sizes.

Now we choose each vertex in V^* independently with probability $p = T^{-1+\varepsilon}$ for some constant $\varepsilon > 0$. Let $V^{**} \subseteq V^*$ be the set of chosen vertices, and let $\mathcal{H}^{**} = (V^{**}, \mathcal{E}_3^{**} \cup \cdots \cup \mathcal{E}_k^{**})$ be the subhypergraph of $\mathcal{H}$ induced on V^{**}. Note that

$$\mathbb{E}\left[|V^{**}|\right] = p|V^*| \ge \frac{N}{k} T^{-1+\varepsilon}. \tag{29}$$

Let $C^{**}(g,i,j)$ and $C^{**}(g,h,i,j)$ be the numbers of 3-cycles and 4-cycles (not containing a 3-cycle) in $\mathcal{H}^{**}$, respectively. Since a 3-cycle covers exactly $(g+h+i-3)$ vertices in a linear hypergraph, inequality (27) yields that for $i \ge 3$,

$$\begin{aligned} \mathbb{E}[C^{**}(g,h,i)] < p^{g+h+i-3} Nk^4 c_g c_h T^{g+h-2} (\ln T)^2 &\le k^4 c_g c_h N T^{-(i-1)+\varepsilon(g+h+i-3)} (\ln T)^2 \\ &< k^4 c_g c_h N T^{-2+3k\varepsilon} (\ln T)^2. \end{aligned}$$

Moreover, since there are no 3-cycles with 3 edges, each of size 2, in $\mathcal{H}$, we have $\mathbb{E}[C^{**}(2,2,2)] = 0$. Hence, we infer

$$\sum_{2 \le g \le h \le i \le k} \mathbb{E}[C^{**}(g,h,i)] < k^7 \max_{2 \le g \le k} \{c_g^2\} \cdot N T^{-2+3k\varepsilon} (\ln T)^2. \tag{30}$$

Similarly, inequality (28) implies that for $j \ge 3$,

$$\begin{aligned} \mathbb{E}[C^{**}(g,h,i,j)] &< p^{g+h+i+j-4} 3Nk^6 c_g c_h c_i T^{g+h+i-3} (\ln T)^3 \\ &= 3k^6 c_g c_h c_i N T^{-(j-1)+\varepsilon(g+h+i+j-4)} (\ln T)^3 \\ &\le 3k^6 \max_{2 \le g \le k} \{c_g^3\} \cdot N T^{-2+4k\varepsilon} (\ln T)^3. \end{aligned}$$

Also since there are no 4-cycles with 4 edges, each of size 2, in $\mathcal{H}$, we have $\mathbb{E}[C^{**}(2,2,2,2)] = 0$. Thus, we infer

$$\sum_{2 \le g \le h \le i \le j \le k} \mathbb{E}[C^{**}(g,h,i,j)] \le 3k^{10} \max_{2 \le g \le k} \{c_g^3\} \cdot N T^{-2+4k\varepsilon} (\ln T)^3. \tag{31}$$

By (26), we have for $i = 2, \ldots, k$ that

$$\mathbb{E}[|\mathcal{E}_i^{**}|] = p^i |\mathcal{E}_i^*| \le p^i k c_i T^{i-1} (\ln T)^{\frac{k-i}{k-1}} \frac{|V^*|}{i} = k c_i T^{-1+\varepsilon i} (\ln T)^{\frac{k-i}{k-1}} \frac{|V^*|}{i}. \tag{32}$$

Chernoff's and Markov's inequalities with (29)–(32) imply that there exists a subhypergraph $\mathcal{H}^{**} = (V^{**}, \mathcal{E}_2^{**} \cup \cdots \cup \mathcal{E}_k^{**})$ induced on V^{**} such that

$$\begin{aligned} |V^{**}| &\ge \frac{1}{2k} N T^{-1+\varepsilon} \\ |\mathcal{E}_i^{**}| &\le (k+2) k c_i T^{-1+\varepsilon i} (\ln T)^{\frac{k-i}{k-1}} \frac{|V^*|}{i} \\ \sum_{2 \le g \le h \le i \le k} C^{**}(g,h,i) &\le (k+2) k^7 \max_{2 \le g \le k} \{c_g^2\} \cdot N T^{-2+3k\varepsilon} (\ln T)^2 \\ \sum_{2 \le g \le h \le i \le j \le k} C^{**}(g,h,i,j) &\le (k+2) 3k^{10} \max_{2 \le g \le k} \{c_g^3\} \cdot N T^{-2+4k\varepsilon} (\ln T)^3. \end{aligned} \tag{33}$$

For $0 < \varepsilon < 1/(4k-1)$ and $T > T_0(k)$, the number of 3-cycles and the number of 4-cycles (not containing a 3-cycle) are much less than $|V^{**}|$.

Let $\mathcal{H}^{***} = (V^{***}, \mathcal{E}_2^{***} \cup \cdots \cup \mathcal{E}_k^{***})$ be the subhypergraph obtained from $\mathcal{H}^{**}$ by removing one vertex from each 3-cycle and 4-cycle (not containing a 3-cycle). With $\varepsilon := 1/(4k)$, we have

$$|V^{***}| \geq \frac{1}{4k} N T^{-1+\varepsilon}. \tag{34}$$

We infer that the average degrees $(t_i^{***})^{i-1}$ of $\mathcal{H}^{***}$ satisfy for $i = 2, \ldots, k$:

$$\begin{aligned}
(t_i^{***})^{i-1} &= \frac{i|\mathcal{E}_i^{***}|}{|V^{***}|} \leq \frac{i|\mathcal{E}_i^{**}|}{|V^{***}|} \overset{(33)}{\leq} (k+2)kc_i T^{-1+\varepsilon i} (\ln T)^{\frac{k-i}{k-1}} \frac{|V^*|}{|V^{***}|} \\
&\leq (k+2)kc_i T^{-1+\varepsilon i} (\ln T)^{\frac{k-i}{k-1}} \frac{N}{|V^{***}|} \\
&\overset{(34)}{\leq} 4k^2(k+2)c_i T^{\varepsilon(i-1)} (\ln T)^{\frac{k-i}{k-1}} \\
&= \frac{4k^2(k+2)}{\varepsilon^{\frac{k-i}{k-1}}} c_i (T^\varepsilon)^{i-1} (\ln(T^\varepsilon))^{\frac{k-i}{k-1}} \\
&\leq 32k^4 c_i (T^\varepsilon)^{i-1} (\ln(T^\varepsilon))^{\frac{k-i}{k-1}}.
\end{aligned}$$

Since the assumption $32k^4 c_i < \frac{1}{16k^2}\binom{k-1}{i-1} 10^{-3\frac{k-i}{k-1}}$ of Lemma 12 is satisfied, this implies that there exists a constant $C_k > 0$ such that

$$\begin{aligned}
\alpha(\mathcal{H}^{***}) &\geq C_k \frac{|V^{***}|}{T^\varepsilon} (\ln(T^\varepsilon))^{\frac{1}{k-1}} \overset{(34)}{\geq} C_k \frac{NT^{-1+\varepsilon}/(4k)}{T^\varepsilon} (\ln(T^\varepsilon))^{\frac{1}{k-1}} \\
&\geq C_k' \frac{N}{T} (\ln T)^{\frac{1}{k-1}},
\end{aligned}$$

which completes our proof of Lemma 15. □

5. Proof of Theorem 6

We are going to show Theorem 6 from Lemma 15 that is a weaker version of Theorem 6. We need to modify two assumptions of Lemma 15: the first assumption is about the upper bounds on t_i^{i-1}, and the second assumption is about the ranges of T and N.

We first change the assumptions on the upper bounds on t_i^{i-1}, and show the following.

Lemma 16. *Let $k \geq 2$ be a fixed integer. Let T and N satisfy $T > T_0(k)$ and $N > N_0(k,T)$. Let $\mathcal{H} = (V, \mathcal{E}_2 \cup \cdots \cup \mathcal{E}_k)$ be a linear hypergraph on N vertices such that there are no 3-cycles and 4-cycles consisting of 2-element edges. Let the average degrees $t_i^{i-1} := i|\mathcal{E}_i|/N$ satisfy that for $i = 2, \ldots, k$*

$$t_i^{i-1} \leq T^{i-1} (\ln T)^{\frac{k-i}{k-1}}. \tag{35}$$

Then, there exists a constant $C_k > 0$ such that

$$\alpha(\mathcal{H}) \geq C_k \frac{N}{T} (\ln T)^{\frac{1}{k-1}} .$$

Proof. We are going to use Lemma 15. To this end, we need to change the assumption (35) to the shape of the assumption (24) in Lemma 15. We have

$$\begin{aligned}
t_i^{i-1} &\leq T^{i-1} (\ln T)^{\frac{k-i}{k-1}} = c_i \cdot \frac{2}{c_i} T^{i-1} \cdot \frac{1}{2} (\ln T)^{\frac{k-i}{k-1}} \\
&\leq c_i \left(\frac{T}{c_i^*}\right)^{i-1} \left(\frac{1}{2} \ln T\right)^{\frac{k-i}{k-1}} \leq c_i \left(\frac{T}{c_i^*}\right)^{i-1} \left(\ln \frac{T}{c_i^*}\right)^{\frac{k-i}{k-1}},
\end{aligned}$$

where $c_i^* := (c_i/2)^{\frac{1}{i-1}}$ and the last inequality holds because T is sufficiently large depending on k.

Now we apply Lemma 15, and we infer

$$\begin{aligned}\alpha(\mathcal{H}) &\geq C_k \frac{N}{T/c_i^*}\left(\ln \frac{T}{c_i^*}\right)^{\frac{1}{k-1}} \geq C_k c_i^* \frac{N}{T}\cdot\frac{1}{2}(\ln T)^{\frac{1}{k-1}} \\ &\geq C_k' \frac{N}{T}(\ln T)^{\frac{1}{k-1}},\end{aligned}$$

where $C_k' := C_k c_i^*/2$ and the second inequality holds since T is sufficiently large depending on k. This completes the proof of Lemma 16. □

In order to show Theorem 6, it only remains to enlarge the ranges of T and N in Lemma 16 as $T \geq 1.5$ and every N.

First, we enlarge the range of T from $T > T_0(k)$ to $T \geq 1.5$. If $1.5 \leq T \leq T_0(k)$, then $T \ln T \geq 1/2$, and hence, Theorem 11 with $t_i^{i-1} \leq (T \ln T)^{i-1}$ implies that

$$\begin{aligned}\alpha(\mathcal{H}) \geq \frac{1}{4}\frac{N}{T\ln T} \geq \frac{1}{4}\frac{N}{T\ln T_0} &= \left(\frac{1}{4(\ln T_0)^{1+\frac{1}{k-1}}}\right)\frac{N}{T}(\ln T_0)^{\frac{1}{k-1}} \\ &\geq \left(\frac{1}{4(\ln T_0)^{1+\frac{1}{k-1}}}\right)\frac{N}{T}(\ln T)^{\frac{1}{k-1}} = C_k\frac{N}{T}(\ln T)^{\frac{1}{k-1}}.\end{aligned}$$

Next, we enlarge the range of N from $N > N_0(k,T)$ to be every N. Let $\mathcal{H} = (V,\mathcal{E})$ be a hypergraph satisfying the assumption in Lemma 16 except for $T > T_0(k)$ and $N > N_0(k,T)$, and suppose that $N \leq N_0(k,T)$ and $T \geq 1.5$.

Let $L > N_0(k,T)/N$, and consider the hypergraph $\mathcal{H}'$ obtained by L vertex-disjoint copies of $\mathcal{H}$. Observe that $\mathcal{H}'$ has LN vertices and its average degree is the same as the average degree of $\mathcal{H}$. Hence $\mathcal{H}'$ satisfies the assumption of Lemma 16, thus

$$\alpha(\mathcal{H}') \geq C_k\frac{LN}{T}(\ln T)^{\frac{1}{k-1}}.$$

Since $\alpha(\mathcal{H}) = \alpha(\mathcal{H}')/L$, we infer

$$\alpha(\mathcal{H}) \geq C_k\frac{N}{T}(\ln T)^{\frac{1}{k-1}},$$

which completes the proof of Theorem 6. □

6. Proof of Theorem 8

It will be convenient here to work with the O-notation. For functions $f, g\colon \mathbb{N} \longrightarrow \mathbb{N}$ and a fixed integer $\ell > 0$, let $f = O_\ell(g)$ mean that there exists a constant $c > 0$, depending only on ℓ, such that $f(n) \leq cg(n)$ for every sufficiently large $n \in \mathbb{N}$.

Proof of the lower bound of Theorem 8. We will use in our arguments some ideas from [2] (compare also [7]). Recall that $\mathcal{H} = \mathcal{H}(n; 2, \ldots, \ell)$ is the hypergraph on the vertex set V, $|V| = n$, in which, for $s = 2, \ldots, \ell$, each s-subset of V is a hyperedge of $\mathcal{H}$, and let a $(u_2, \ldots u_\ell)$-bounded edge-coloring Δ of $\mathcal{H}$ be given. We define another hypergraph $\mathcal{G} = (V, E)$ as follows: If e_1 and e_2 are hyperedges in $\mathcal{H}$ with the same color in c, then $e_1 \cup e_2 \in E(\mathcal{G})$. Let E_{2i} be the set of hyperedges of $\mathcal{G}$ of size $2i$, hence $E = \bigcup_{i=2}^{\ell} E_{2i}$. Observe that if $I \subset V$ is an independent set of $\mathcal{G}$, then the subhypergraph $\mathcal{H}[I]$ of $\mathcal{H}$ induced on I is totally multicolored. Hence, a lower bound on $f_\Delta(n; u_2, \ldots, u_\ell)$ can be obtained by finding an independent set in $\mathcal{G}$. Therefore, it suffices to show the following:

(i) Under assumption (6), we have

$$\alpha(\mathcal{G}) \geq c_1 \left(\frac{n^s}{u_s}\right)^{\frac{1}{2s-1}}.$$

(ii) Under assumptions (8) and (9), we have

$$\alpha(\mathcal{G}) \geq c_3 \left(\frac{n^s}{u_s}\ln n\right)^{\frac{1}{2s-1}}.$$

We first prove (i). Let assumption (6) hold. Set

$$p = \left(\frac{1}{n^{s-1}u_s}\right)^{\frac{1}{2s-1}}.$$

Let R be a random subset of V obtained by choosing each vertex independently with probability p. Note that with high probability $|R| = np(1+o(1))$.

For $i = 2, \ldots, \ell$, let E_{2i}^R be the set of all $2i$-element hyperedges in the subhypergraph $\mathcal{G}[R]$ induced on R. To estimate the expected numbers $\mathbb{E}(|E_{2i}^R|)$ of $2i$-element hyperedges, choose an i-element hyperedge e in $\binom{n}{i}$ ways. Less than u_i other hyperedges have the same color as e, thus

$$|E_{2i}| \leq \frac{n^i u_i}{2(i!)} \qquad \text{hence,} \qquad \mathbb{E}(|E_{2i}^R|) \leq \frac{n^i u_i p^{2i}}{2(i!)}.$$

Consequently, Markov's inequality gives for $i = 2, \ldots, \ell$ that

$$\Pr\left[|E_{2i}^R| > \ell \cdot \frac{n^i u_i p^{2i}}{2(i!)}\right] \leq \frac{1}{\ell}.$$

Therefore, there exists a vertex set $R \subset V$ such that

- $|R| \geq np/2$,
- $|E_{2i}^R| \leq \ell \cdot \dfrac{n^i u_i p^{2i}}{2(i!)}$ for $i = 2, \ldots, \ell$.

The average degree t_{2i}^{2i-1} for the $2i$-element hyperedges in the subhypergraph $\mathcal{G}[R]$ satisfies

$$t_{2i}^{2i-1} \leq \left(2i\ell \cdot \frac{n^i u_i p^{2i}}{2(i!)}\right) \Big/ (np/2) = \frac{2\ell}{(i-1)!} n^{i-1} u_i p^{2i-1},$$

and hence, for $i = 2, \ldots, \ell$, and some constant $c_\ell > 0$, it is

$$t_{2i} \leq c_\ell \left(n^{i-1}u_i\right)^{1/(2i-1)} p \overset{(6)}{\leq} c_\ell \left(n^{s-1}u_s\right)^{1/(2s-1)} p.$$

By Theorem 11 we infer that

$$\alpha(\mathcal{G}) \geq \frac{1}{4} \frac{np/2}{c_\ell \left(n^{s-1}u_s\right)^{1/(2s-1)} p} = c_1' \left(\frac{n^s}{u_s}\right)^{1/(2s-1)},$$

for some constant $c_1' > 0$, depending only on ℓ, which completes the proof of (i).

Next we show (ii). Suppose that (8) and (9) hold. Set

$$p = \left(\frac{1}{n^{s-1}u_s}\right)^{\frac{1}{2s-1}} \omega, \qquad \text{where } \omega := \left(u_s^2/n\right)^{\frac{1}{2(2s-1)(2\ell+1)}}. \tag{36}$$

As above, let R be a random subset of V obtained by choosing each vertex independently with probability p. We again have that

- $|R| = np(1+o(1))$ with high probability,

- For $i = 2, \ldots, \ell$,

$$\Pr\left[|E_{2i}^R| > 3\ell \cdot \frac{n^i u_i p^{2i}}{2(i!)}\right] \leq \frac{1}{3\ell}.$$

Next, we consider 2-cycles in $\mathcal{G}$. For integers $2 \leq i, j, k \leq \ell$, let $C(2i, 2j, k)$ be the family of all 2-cycles which consist of two distinct hyperedges in $\mathcal{G}$ with one of size $2i$ and the other of size $2j$ sharing exactly k vertices. Note that the condition on k is either $2 \leq k \leq 2i-1$ for $i = j$ or $2 \leq k \leq 2i$ for $i < j$. We estimate $|C(2i, 2j, k)|$ as follows. Fix the first hyperedge $e \in E_{2i}$ as an arbitrary one in at most $n^i u_i$ ways. Let $e_1 \cup e_2 \in E_{2j}$ ($e_1, e_2 \in E_j(\mathcal{H})$) be the second hyperedge in $\mathcal{G}$ and let $k_1 = |e_1 \cap e|$ and $k_2 = |e_2 \cap e|$. Without loss of generality, we assume $k_1 \geq k_2$.

- Suppose $k_1 = k$ and $k_2 = 0$. The number of choices of e_1 is at most $O_\ell(n^{j-k})$. Since the color of e_1 is determined, the number of choices of e_2 is at most $u_j \leq n$ because of the assumption that each color class is a matching. Hence, the number of choices of (e_1, e_2) is at most $O_\ell(n^{j-k+1})$.
- Otherwise, we have $k_2 \neq 0$. The number of choices of e_1 is at most $O_\ell(n^{j-k_1})$. Then, the number of choices of e_2 is at most $O_\ell(1)$. Hence, the number of choices of (e_1, e_2) is at most $O_\ell(n^{j-k_1})$. By minimizing k_1, we have the upper bound $O_\ell\left(n^{j-\lceil k/2 \rceil}\right)$.

Consequently, we have that for all $k \geq 2$,

$$|C(2i, 2j, k)| = O_\ell\left(|E_{2i}| \cdot n^{j-\lceil k/2 \rceil}\right) = O_\ell\left(u_i n^{i+j-\lceil k/2 \rceil}\right).$$

Let $C^R(2i, 2j, k)$ be the random set of all 2-cycles in $C(2i, 2j, k)$ that are contained in R. Let

$$\begin{aligned} C^R(2i, 2i) &= \bigcup_{k=2}^{2i-1} C^R(2i, 2i, k) \quad \text{for } 2 \leq i \leq \ell, \qquad \text{and} \\ C^R(2i, 2j) &= \bigcup_{k=2}^{2i} C^R(2i, 2j, k) \quad \text{for } 2 \leq i < j \leq \ell. \end{aligned}$$

Since $\mathbb{E}\left[|C^R(2i, 2j, k)|\right] = |C(2i, 2j, k)| p^{2i+2j-k}$, we infer that for $2 \leq i \leq \ell$,

$$\begin{aligned} \mathbb{E}\left[|C^R(2i, 2i)|\right] &= \sum_{k=2}^{2i-1} \mathbb{E}\left[|C^R(2i, 2i, k)|\right] \overset{(np \gg 1)}{=} O_\ell\left(u_i n^{2i-1} p^{4i-2} + u_i n^{i+1} p^{2i+2}\right) \\ &\overset{(np^2 \ll 1)}{=} O_\ell\left(u_i n^{i+1} p^{2i+2}\right), \end{aligned}$$

and that for $2 \leq i < j \leq \ell$,

$$\begin{aligned} \mathbb{E}\left[|C^R(2i, 2j)|\right] &= \sum_{k=2}^{2i} \mathbb{E}\left[|C^R(2i, 2j, k)|\right] \overset{(np \gg 1)}{=} O_\ell\left(u_i n^{j+i-1} p^{2j+2i-2} + u_i n^j p^{2j}\right) \\ &\overset{(np^2 \ll 1)}{=} O_\ell\left(u_i n^j p^{2j}\right). \end{aligned}$$

Using Markov's inequality, it simultaneously holds with probability bigger than $2/3$ that

$$\begin{aligned} |C^R(2i, 2i)| &= O_\ell\left(u_i n^{i+1} p^{2i+2}\right) \quad \text{for } 2 \leq i \leq \ell, \qquad \text{and} \\ |C^R(2i, 2j)| &= O_\ell\left(u_i n^j p^{2j}\right) \quad \text{for } 2 \leq i < j \leq \ell. \end{aligned}$$

Therefore, there exists a subset $R \subset V$ such that for some constant $c > 0$, depending only on ℓ, we have that

- $|R| \geq np/2$
- $|E_{2i}^R| \leq c u_i n^i p^{2i}$ for $2 \leq i \leq \ell$

- $|C^R(2i,2i)| \le cu_i n^{i+1}p^{2i+2}$ for $2 \le i \le \ell$, and
  $|C^R(2i,2j)| \le cu_i n^j p^{2j}$ for $2 \le i < j \le \ell$.

We claim that $\sum_{i=2}^{\ell} |C^R(2i,2i)| \ll np$ and $\sum_{2\le i<j\le \ell} |C^R(2i,2j)| \ll np$. For the first inequality, it suffices to check that for $2 \le i \le \ell$,

$$u_i n^{i+1}p^{2i+2} \ll np, \quad \text{that is,} \quad u_i p(np^2)^i \ll 1,$$

indeed, we have that

$$u_i p \overset{(8)}{\le} \left(\frac{u_s^2}{n}\right)^{\frac{i-1}{2s-1}} \omega (\ln n)^{\frac{2(s-i)}{2s-1}} \quad \text{and} \quad np^2 = \left(\frac{n}{u_s^2}\right)^{\frac{1}{2s-1}} \omega^2. \tag{37}$$

Hence,

$$u_i p(np^2)^i \le \left(\frac{n}{u_s^2}\right)^{\frac{1}{2s-1}} \omega^{2i+1} (\ln n)^{\frac{2(s-i)}{2s-1}} \ll 1,$$

where the last inequality follows from $u_s \ge n^{\frac{1}{2}+\varepsilon}$ and $\omega = \left(u_s^2/n\right)^{\frac{1}{2(2s-1)(2\ell+1)}}$. Next, for the second inequality, similarly it follows that $u_i n^j p^{2j} \ll np$ for $2 \le i < j \le \ell$.

After deleting a vertex from each member of $C^R(2i,2i)$ or $C^R(2i,2j)$, there exists a vertex set $U \subset R$ such that

- $|U| \ge \dfrac{np}{4}$
- $|E_{2i}^U| \le cu_i n^i p^{2i}$ for $2 \le i \le \ell$.
- There is no 2-cycle in the subhypergraph $\mathcal{G}[U]$ induced on U.

Set

$$T \;:=\; c^* p (n^{s-1}u_s)^{\frac{1}{2s-1}} (\ln n)^{\frac{2s-2\ell}{(2s-1)(2\ell-1)}}, \tag{38}$$

where $c^* > 0$ is a suitable large constant to be fixed later. We now check that the average degree t_{2i}^{2i-1} of the subhypergraph $\mathcal{G}[U]$ with hyperedges of size $2i$ satisfies (4), that is,

$$t_{2i}^{2i-1} \le T^{2i-1} \cdot (\ln T)^{\frac{2\ell-2i}{2\ell-1}}. \tag{39}$$

First, observe that

$$t_{2i}^{2i-1} \le 8\ell c u_i n^{i-1} p^{2i-1} = 8\ell c (u_i p)(np^2)^{i-1} \overset{(37)}{=} 8\ell c \omega^{2i-1} (\ln n)^{\frac{2(s-i)}{2s-1}}.$$

On the other hand, since $u_s \ge n^{1/2+\varepsilon}$, we have $\ln T \ge c' \ln n$ for some constant $c' = c'(\ell,\varepsilon) > 0$, and hence,

$$T^{2i-1} \cdot (\ln T)^{\frac{2\ell-2i}{2\ell-1}} \ge (c^*)^{2i-1} (c')^{\frac{2\ell-2i}{2\ell-1}} \omega^{2i-1} (\ln n)^{\frac{2(s-i)}{2s-1}}.$$

With a suitable large choice of $c^* = c^*(\ell,\varepsilon) > 0$, depending on c and c', we infer (39).

Theorem 6 implies that there exist positive constant c_ℓ, depending only on ℓ, such that

$$\alpha(\mathcal{G}) \ge c_\ell \frac{np}{T} (\ln T)^{\frac{1}{2\ell-1}} = c_3 \left(\frac{n^s}{u_s} \ln n\right)^{\frac{1}{2s-1}},$$

where $c_3 > 0$ is a constant, depending only on ℓ and ε, which completes the proof of (ii). □

Proof of the upper bound of Theorem 8. We will show the following using some ideas from [2] and [3]: For each integer k with $2 \le k \le \ell$, there exists a positive constant $C = C(k)$ such that for every sufficiently large n,

$$f(n; u_2, ..., u_\ell) \le C \left(\frac{n^k}{u_k} \ln n\right)^{\frac{1}{2k-1}}. \tag{40}$$

This with $k = s$ implies the upper bound in Theorem 8.

For a proof of (40), let $\mathcal{H}_k$ be the subhypergraph of $\mathcal{H}$ on the vertex set V only with all k-element hyperedges. We define a random edge-coloring of $\mathcal{H}_k$ as follows. Set

$$m = c_0 \frac{n^k}{u_k},$$

where c_0 is a constant with $0 \leq c_0 \leq 1/(8e^2(k!))$. (We ignore divisibility constraints in our arguments, as there is enough room in the calculations.) Let $M_1, \ldots, M_m$ be random matchings chosen uniformly and independently from the set of all matchings of size u_k on V, and let $U_0 = \emptyset$ and $U_i = \bigcup_{j \leq i} M_j$ for $i = 1, \ldots, m$. We color all hyperedges in $M_i \setminus U_{i-1}$ by color i, and color the remaining ones with distinct new colors.

In order to prove (40), it suffices to show that, for $x = C\left(\frac{n^k}{u_k} \ln n\right)^{\frac{1}{2k-1}}$, where $C > 0$ is a suitable constant to be fixed later, we have

$$\Pr\left[\exists X \subset V \text{ such that } |X| = x \text{ and } X \text{ is totally multicolored } \right] = o(1).$$

For its proof, let $X \subseteq V$ be an arbitrarily fixed subset of V of size x. We will show that X is totally multicolored with very small probability. For $i = 1, \ldots, m$, let Y_i be the number of pairs $\{S, T\}$ of hyperedges in $M_i \setminus U_{i-1}$ contained in X. Observe that X is totally multicolored if and only if simultaneously $Y_i = 0$ for $i = 1, \ldots, m$. We can show that $Y_i = 0,\ i = 0, \ldots, m$, hold with very small probability under the condition that the intersection size $|U_m \cap [X]^k|$ is small, where $[X]^k$ denotes the family of all k-element hyperedges in X. To this end, let A be the event that $|U_m \cap [X]^k| \leq c_1 x^k$ where $c_1 = 1/(4(k!))$. We have the following:

$$\begin{aligned} \Pr\left[X \text{ is totally multicolored } \right] &= \Pr\left[Y_1 = 0, \ldots, Y_m = 0\right] \\ &\leq \Pr[A^c] + \Pr\left[Y_1 = 0, \ldots, Y_m = 0, A\right] \\ &\leq \Pr[A^c] + \Pr\left[Y_1 = 0, \ldots, Y_m = 0 \mid A\right]. \end{aligned} \tag{41}$$

We will use the following two claims:

Claim 17. *For every sufficiently large n,*

$$\Pr\left[A^c\right] \leq \exp\left(-c_1 x^k\right). \tag{42}$$

Claim 18. *For every sufficiently large n,*

$$\Pr\left[Y_1 = 0, \ldots, Y_m = 0 \mid A\right] \leq \exp\left(-\frac{c_0 c_1^2}{4} \frac{u_k x^{2k}}{n^k}\right). \tag{43}$$

The union bound and Claims 17 and 18 together with (41) yield that

$$\begin{aligned} & \Pr\left[\exists X \subset V \text{ such that } |X| = x \text{ and } X \text{ totally multicolored}\right] \\ \leq\ & \binom{n}{x}\left(\exp\left(-\frac{c_0 c_1^2}{4} \frac{u_k x^{2k}}{n^k}\right) + \exp\left(-c_1 x^k\right)\right) \\ \leq\ & \exp\left(x \ln n\right) \cdot \left(\exp\left(-\frac{c_0 c_1^2}{4} \frac{u_k x^{2k}}{n^k}\right) + \exp\left(-c_1 x^k\right)\right). \end{aligned} \tag{44}$$

By choosing the constant $C > (4/(c_0 c_1^2))^{\frac{1}{2k-1}}$, the term (44) goes to 0 as n tends to ∞, which completes our proof of (6). It remains to prove Claims 17 and 18.

First, we prove Claim 17. Since $|U_m \cap [X]^k| \leq \sum_{i=1}^m |M_i \cap [X]^k|$ and the events $|M_i \cap [X]^k| \geq t_i$, $i = 1, \ldots, m$, are independent, we infer that

$$\Pr\left[|U_m \cap [X]^k| > t\right] \leq \Pr\left[\sum_{i=1}^m |M_i \cap [X]^k| \geq t\right] \leq \sum_{\substack{(t_i)_{i=1}^m \text{ s.t.} \\ t_i \geq 0, \sum_{i=1}^m t_i = t}} \prod_{i=1}^m \Pr\left[|M_i \cap [X]^k| \geq t_i\right].$$

Now we estimate $\Pr\left[|M_i \cap [X]^k| \geq t_i\right]$ for integers $1 \leq i \leq m$ and $t_i \geq 0$. There are $\binom{u_k}{t_i}$ choices for selecting t_i hyperedges in M_i. Then the t_i hyperedges are contained in X with probability $\binom{x}{kt_i}/\binom{n}{kt_i}$, hence

$$\Pr\left[|M_i \cap [X]^k| \geq t_i\right] \leq \binom{u_k}{t_i} \frac{\binom{x}{kt_i}}{\binom{n}{kt_i}} \leq \left(\frac{u_k x^k}{n^k}\right)^{t_i}. \tag{45}$$

Thus, we infer that

$$\begin{aligned}
\Pr\left[|U_m \cap [X]^k| > t\right] &\leq \sum_{\substack{(t_i)_{i=1}^m \text{ s.t.} \\ t_i \geq 0, \sum_{i=1}^m t_i = t}} \prod_{i=1}^m \left(\frac{u_k x^k}{n^k}\right)^{t_i} \leq \binom{t+m-1}{t} \left(\frac{u_k x^k}{n^k}\right)^t \\
&\leq \left(\frac{e(t+m)}{t}\right)^t \left(\frac{u_k x^k}{n^k}\right)^t \leq \left(\frac{e(t+m)u_k x^k}{tn^k}\right)^t.
\end{aligned}$$

Take $t = c_1 x^k$ and note that $t = o(m)$. Consequently,

$$\Pr\left[|U_m \cap [X]^k| > c_1 x^k\right] \leq \left(\frac{2emu_k x^k}{c_1 x^k n^k}\right)^{c_1 x^k} = \left(\frac{2ec_0}{c_1}\right)^{c_1 x^k} \leq e^{-c_1 x^k},$$

where the last inequality follows from $0 \leq c_0 \leq 1/(8e^2(k!))$ and $c_1 = 1/(4(k!))$. This completes our proof of Claim 17.

Next, we prove Claim 18. First, we have that

$$\Pr\left[Y_1 = 0, \ldots, Y_m = 0 \;\middle|\; A\right] = \prod_{i=1}^m \Pr\left[Y_i = 0 \;\middle|\; A, Y_j = 0 \text{ for } 1 \leq j \leq i-1\right]. \tag{46}$$

For simplification, let B_i denote the event that A happens and $Y_j = 0$ for $1 \leq j \leq i$. Next we upper bound $\Pr[Y_i = 0 \mid B_{i-1}]$, or equivalently, we lower bound $\Pr[Y_i \geq 1 \mid B_{i-1}]$.

To this end, it is useful to consider $\mathbb{E}[Y_i \mid B_{i-1}]$. The condition that A holds implies $|U_{i-1} \cap [X]^k| \leq c_1 x^k$, that is,

$$|[X]^k \setminus U_{i-1}| \geq \binom{x}{k} - c_1 x^k \geq \left(\frac{1}{2(k!)} - c_1\right) x^k = c_1 x^k.$$

For each hyperedge $S \in [X]^k$ there are at most $k\binom{x-1}{k-1}$ hyperedges in X which are not disjoint from S. Hence, the number of pairs $\{S, T\} \in [[X]^k \setminus U_{i-1}]^2$ of hyperedges with $S \cap T = \emptyset$ is, for every sufficiently large n, at least

$$\frac{1}{2} c_1 x^k \left(c_1 x^k - k\binom{x-1}{k-1}\right) > \frac{c_1^2 x^{2k}}{3}.$$

For disjoint hyperedges S and T,

$$\Pr\left[S, T \in M_i\right] = \frac{u_k(u_k - 1)}{\binom{n}{k}\binom{n-k}{k}} \geq \frac{u_k^2}{n^{2k}},$$

and therefore

$$\mathbb{E}[Y_i \mid B_{i-1}] \;>\; \frac{c_1^2}{3}\Big(\frac{u_k x^k}{n^k}\Big)^2. \tag{47}$$

Now we estimate $\Pr[Y_i \geq 1 \mid B_{i-1}]$ by using $\mathbb{E}[Y_i \mid B_{i-1}]$. We have that

$$\begin{aligned}\Pr[Y_i \geq 1 \mid B_{i-1}] &= \mathbb{E}[Y_i \mid B_{i-1}] - \sum_{j\geq 2}(j-1)\Pr[Y_i = j \mid B_{i-1}] \\ &= \mathbb{E}[Y_i \mid B_{i-1}] - \sum_{j\geq 2}\Pr[Y_i \geq j \mid B_{i-1}]. \end{aligned}\tag{48}$$

Observe that for j pairwise distinct two-element sets, the underlying set has cardinality at least $\lceil \sqrt{2j+1}\,\rceil$. Hence,

$$\Pr[Y_i \geq j \mid B_{i-1}] \leq \Pr\Big[|M_i \cap [X]^k| \geq \lceil \sqrt{2j+1}\,\rceil\Big] \overset{(45)}{\leq} \left(\frac{u_k x^k}{n^k}\right)^{\lceil \sqrt{2j+1}\,\rceil}. \tag{49}$$

Consequently, it follows from (47)–(49) and $x^k = o\left(n^k/u_k\right)$ that

$$\Pr[Y_i \geq 1 \mid B_{i-1}] \geq \frac{c_1^2}{4}\left(\frac{u_k x^k}{n^k}\right)^2,$$

that is,

$$\Pr[Y_i = 0 \mid B_{i-1}] \leq 1 - \frac{c_1^2}{4}\left(\frac{u_k x^k}{n^k}\right)^2.$$

Therefore, (46) gives that

$$\Pr\Big[Y_1 = 0, \ldots, Y_m = 0 \;\Big|\; A\Big] \;\leq\; \left(1 - \frac{c_1^2}{4}\frac{u_k^2 x^{2k}}{n^{2k}}\right)^m \leq \exp\left(-\frac{c_1^2}{4}\frac{u_k^2 x^{2k} m}{n^{2k}}\right) \leq \exp\left(-\frac{c_0 c_1^2}{4}\frac{u_k x^{2k}}{n^k}\right),$$

which completes our proof of Claim 18. □

References

[1] M. Ajtai, J. Komlós, J. Pintz, J. Spencer and E. Szemerédi, *Extremal Uncrowded Hypergraphs*, Journal of Combinatorial Theory Ser. A 32, 1982, 321–335. 1, 4, 1, 3, 13

[2] N. Alon, H. Lefmann and V. Rödl, *On an Anti-Ramsey Type Result,* Coll. Math. Soc. János Bolyai, 60. Sets, Graphs and Numbers, 1991, 9–22. 1, 6, 6

[3] L. Babai, *An Anti-Ramsey Theorem,* Graphs and Combinatorics 1, 1985, 23–28. 1, 6

[4] C. Bertram-Kretzberg and H. Lefmann, *The Algorithmic Aspects of Uncrowded Hypergraphs,* SIAM Journal on Computing 29, 1999, 201–230. 1

[5] R. A. Duke, H. Lefmann and V. Rödl, *On Uncrowded Hypergraphs,* Random Structures & Algorithms 6, 1995, 209–212. 1, 5

[6] A. Fundia, *Derandomizing Chebychev's Inequality to Find Independent Sets in Uncrowded Hypergraphs*, Random Structures & Algorithms 8, 1996, 131–147. 1, 3

[7] H. Lefmann, V. Rödl and B. Wysocka, *Multicolored Subsets in Colored Hypergraphs,* Journal of Combinatorial Theory Series A 74, 1996, 209–248. 1, 6

[8] P. Turán, *On an Extremal Problem in Graph Theory,* Mat. Fiz. Lapok 48, 1941, 436–452. 1, 1

[9] J. Spencer, *Turán's Theorem for k-Graphs*, Discrete Mathematics 2, 1972, 183–186. 1, 1, 2, 11

Department of Mathematics, Duksung Women's University, Seoul 01369, South Korea
E-mail address: sanglee242@duksung.ac.kr, sjlee242@gmail.com

Fakultät für Informatik, TU Chemnitz, D-09107 Chemnitz, Germany
E-mail address: lefmann@informatik.tu-chemnitz.de